

A Computational Model for CO₂-Dependence on Temperature in the Vostok Ice cores

[latexpage]


The [Vostok Ice core provides](#) a more than 400000 year view into the climate history with several cycles between ice ages and warm periods.

It has become clear that [CO₂ data](#) are lagging [temperature data](#) by several centuries. One difficulty arises from the necessity that CO₂ is measured in the gas bubbles whereas temperature is determined from a deuterium proxy in the ice. Therefore there is a different way of determining the age for the two parameters – for CO₂ there is a „gas age“, whereas the temperature series is assigned an „ice age“. There are estimates of how much older the „ice age“ is in comparison to the gas age. But there is uncertainty, so we will have to tune the relation between the two time scales.


Preprocessing the Vostok data sets

In order to perform model based computations with the two data sets, the original data must be converted into equally spacially sampled data sets. This is done by means of linear interpolation. The sampling interval is chosen 100 years, which is approximately the sampling interval of the temperature data. Apart from this, the data sets must be reversed, and the sign of the time axis must be set to negative values.


Here is the re-sampled temperature data set from -370000 years to -10000 years overlaid over the original temperature data:



And here the corresponding CO₂-data set:



The two data sets are now superimposed:



Data model

Due to the fact of the [very good predictive value of the temperature dependent sink model for current emission, concentration, and temperature data](#) (equation 2) , we will use the same model based on CO₂ mass balance, and possible linear dependence of CO₂ changes on concentration and temperature, but obviously without the anthropogenic emissions. Also the time interval is no longer a single year, but a century.

G_i is growth of CO₂-concentration C_i during century i :

$$G_i = C_{i+1} - C_i$$

T_i is the average temperature during century i . The [model equation without anthropogenic emissions is](#):

$$G_i = x_1 \cdot C_i + x_2 \cdot T_i + \text{const}$$

After estimating the 3 parameters x_1 , x_2 , and const from G_i , C_i , and T_i by means of ordinary least Squares,


the modelled CO₂ data \hat{C}_i are recursively reconstructed by means of the model, the first actual concentration value of the data sequence C_0 , and the temperature data:

$$\hat{C}_0 = C_0$$

$$\hat{C}_{i+1} = \hat{C}_i - x_1 \cdot \hat{C}_i - x_2 \cdot T_i - \text{const}$$

Results – reconstructed CO₂ data

The standard deviation of $\{\hat{C}_i - C_i\}$ measures the quality of the reconstruction. Minimizing this standard deviation by shifting the temperature data is optimized, when the temperature data is shifted 1450..1500 years to the past:



Here are the corresponding estimated model parameters and the statistical quality measures from the Python OLS package:

```


-----
Model:                OLS                Adj. R-squared:      0.108
Dependent Variable:  y                   AIC:                8772.4952
Date:                2024-02-28 11:26    BIC:                8791.0613
No. Observations:   3600                Log-Likelihood:     -4383.2
Df Model:            2                   F-statistic:        219.7
Df Residuals:        3597                Prob (F-statistic): 9.67e-91
R-squared:           0.109                Scale:              0.66904
-----

```


	Coef.	Std.Err.	t	P> t	[0.025	0.975]
x1	0.0133	0.0009	15.3030	0.0000	0.0116	0.0149
x2	-0.1799	0.0086	-20.8927	0.0000	-0.1968	-0.1630
const	-3.8965	0.2298	-16.9525	0.0000	-4.3471	-3.4458

The interpretation is, that there is a carbon sink of 1.3% per century, and an emission increase of 0.18 ppm per century and 1 degree temperature increase.

Modelling the sinks (-G\$_i\$) results in this diagram:



And the main result, the reconstruction of C0\$_2\$ data from the temperature extended sink model looks quite remarkable:



Equilibrium Relations

The equilibrium states are more meaningful than the incremental changes. The equilibrium is defined by equality of CO₂ sources and sinks, resulting in $G_i = 0$. This creates a linear relation between CO₂ concentration C and Temperature T:

$$C = \frac{0.1799 \cdot T + 3.8965}{0.0133} \text{ ppm}$$

For the temperature anomaly $T=0$ we therefore get the CO₂ concentration of

$$C_{T=0} = \frac{3.8965}{0.0133} \text{ ppm} = 293 \text{ ppm}.$$

The difference of this to the modern data can be explained by different temperature references. Both levels are remarkably close, considering the very different environmental conditions.

And relative change is

$$\frac{dC}{dT} = 13.5 \frac{\text{ppm}}{^{\circ}\text{C}}$$

This is considerably different from the modern data, where we got \$ $66.5 \frac{\text{ppm}}{^{\circ}\text{C}}$ \$.

There is no immediate explanation for this deviation. We need, however, consider the fact that we have time scale differences of at least 100 if not more. Therefore we can expect totally different mechanisms at work.